

Novartis Entresto® granted expanded indication in chronic heart failure by FDA

Feb 16, 2021

- Entresto is the first and only therapy approved in the US to treat patients diagnosed with guideline-defined heart failure to include both those with heart failure with reduced ejection fraction (HFrEF) and many with heart failure with preserved ejection fraction (HFpEF)(1-3)
- Expanded indication enables potential treatment of more adults with left ventricular ejection fraction (LVEF) below normal, the group where benefits are most clearly evident(1)
- Of the more than 6 million Americans suffering from chronic heart failure (CHF), approximately 5 million may be appropriate for treatment with Entresto(3,4)

EAST HANOVER, N.J., Feb. 16, 2021 -- Novartis today announced that the US Food and Drug Administration (FDA) has approved the following expanded indication for Entresto® (sacubitril/valsartan): to reduce the risk of cardiovascular death and hospitalization for heart failure in adult patients with chronic heart failure.¹ Benefits are most clearly evident in patients with left ventricular ejection fraction (LVEF) below normal.¹ The label also states LVEF is a variable measure and clinical judgment should be used in deciding whom to treat.¹

Experience the interactive Multichannel News Release here:

<https://www.multivu.com/players/English/8848351-novartis-entresto-fda-approval/>

For the first time, there is a treatment with benefit for patients diagnosed with guideline-defined heart failure that includes both those with heart failure with reduced ejection fraction (HFrEF) and many with heart failure with preserved ejection fraction (HFpEF).¹⁻³

"This approval is a significant advancement, providing a treatment to many patients who were not eligible for treatment before because their ejection fraction was above the region we normally considered reduced. Until now, treatment for these patients was largely empiric," said Scott Solomon, MD, Professor of Medicine at Harvard Medical School and Brigham and Women's Hospital, and PARAGON-HF Executive Committee Co-Chair. "We can now offer a treatment to a wider range of patients who have an LVEF below normal."

This label expansion is based on efficacy and safety evidence observed in PARAGON-HF, the largest and only Phase III active-controlled study to date in patients with guideline-defined HFpEF.^{2,5,6} The greatest benefit was shown in patients with LVEF below normal.⁶

Approximately 6 million Americans are living with chronic heart failure (CHF).⁴ Approximately 3 million have HFrEF, and of the remaining 3 million, about 2 million have HFpEF with LVEF below normal.²⁻⁴ The prevalence of heart failure (HF) is increasing as the population ages.⁴ Patients often face worsening symptoms that result in frequent HF hospitalizations.⁷ Each hospitalization event is associated with worsening long-term prognosis.⁷ Approximately one in four patients are re-admitted for HF and 10 percent may die within 30 days of discharge.^{8,9} Overall CHF death rates remain significantly high, with up to half of patients dying within five years of a HF diagnosis.⁴

"We are proud of our goal to reimagine medicine. This commitment has enabled us to bring Entresto to millions more heart failure patients in the US, many of whom did not have an approved treatment option until now," said Marie-France Tschudin, President, Novartis Pharmaceuticals. "This achievement would not have been possible without tremendous dedication from investigators, patients in our clinical trials and the advocacy community, to whom we are extremely grateful."

About our longstanding commitment to heart failure

Our goal is to reimagine medicine for heart failure patients. Novartis established the largest global clinical program in the HF disease area across the pharma industry to date. Known as FortiHFy, it is comprised of more than 40 clinical studies designed to generate an array of additional data on efficacy, quality of life, patient-reported outcomes and real-world evidence with Entresto, as well as to extend understanding of heart failure. FortiHFy includes trials across the spectrum of heart failure, such as PARADIGM-HF, PIONEER-HF, TRANSITION, PROVE-HF, PARAGON-HF and PARAMOUNT. Worldwide, it is estimated that more than 30,000 patients have participated in the Entresto clinical trials program, and it is estimated that more than 2.8 million patients are on treatment with Entresto today.

About Entresto

Entresto (sacubitril/valsartan) is a prescription medicine to treat adults with long-lasting (chronic) heart failure to help reduce the risk of death and hospitalization. Entresto works better when the heart cannot pump a normal amount of blood to the body.¹ Entresto is a twice-a-day prescription medicine that works by enhancing the beneficial neurohormonal systems (natriuretic peptide system) while simultaneously inhibiting the harmful effects of the overactive renin-angiotensin-aldosterone system (RAAS).^{1,10} Some other heart failure medicines only block the harmful effects of the overactive RAAS. Entresto contains the neprilysin inhibitor sacubitril and the ARB valsartan.¹ Entresto film-coated tablets are available in three dosage strengths: 24/26 mg, 49/51 mg and 97/103 mg (sacubitril/valsartan).¹ These doses are referred to as 50 mg, 100 mg and 200 mg in the clinical trial literature including The New England Journal of Medicine publication of the results of PARADIGM-HF.¹ In adult patients, the target maintenance dose of Entresto is 97/103 mg twice daily as tolerated by the patient.¹

IMPORTANT SAFETY INFORMATION

Entresto can harm or cause death to an unborn baby. Patients should talk to their doctor about other ways to treat heart failure if they plan to become pregnant. If a patient gets pregnant during treatment with Entresto, she should tell her doctor right away.

Patients should not take Entresto if they are allergic to any of the ingredients in Entresto; have had an allergic reaction including swelling of the face, lips, tongue, throat or trouble breathing while taking a type of medicine called an angiotensin-converting enzyme (ACE) inhibitor or angiotensin II receptor blocker (ARB); or take an ACE inhibitor medicine. Patients should not take Entresto for at least 36 hours before or after they take an ACE inhibitor medicine. Patients should talk with their doctor or pharmacist before taking Entresto if they are not sure if they take an ACE inhibitor medicine. Patients should not take Entresto if they have diabetes and take a medicine that contains aliskiren.

Before they take Entresto, patients should tell their doctor about all of their medical conditions, including if they have a history of hereditary angioedema; have kidney or liver problems; are pregnant or plan to become pregnant; are breastfeeding or plan to breastfeed. Patients should either take Entresto or breastfeed. They should not do both.

Patients should tell their doctor about all the medicines they take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. They should especially tell their doctor if they take potassium supplements or a salt substitute; nonsteroidal anti-inflammatory drugs (NSAIDs); lithium; or other medicines for high blood pressure or heart problems such as an ACE inhibitor, ARB, or aliskiren.

Entresto may cause serious side effects including serious allergic reactions causing swelling of the face, lips, tongue, and throat (angioedema) that may cause trouble breathing and death. Patients should get emergency medical help right away if they have symptoms of angioedema or trouble breathing. Patients are not to take Entresto again if they have had angioedema while taking Entresto. People who are black or who have had angioedema may have a higher risk of having angioedema if they take Entresto. Entresto may cause low blood pressure (hypotension), which may be more common if you take water pills. Patients are to call their doctor if they become dizzy or lightheaded, or they develop extreme fatigue. Entresto may cause kidney problems or an increased amount of potassium in the blood (hyperkalemia).

The most common side effects in adults were low blood pressure, high potassium, cough, dizziness, and kidney problems.

Please see full Prescribing Information, including Boxed WARNING available at <https://www.novartis.us/sites/www.novartis.us/files/entresto.pdf>

Patients are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch, or call 1-800-FDA-1088.

Novartis is committed to providing patients with affordable access and resources through Entresto Central. For more information, please call 1-888-ENTRESTO or visit www.entresto.com.

About Novartis

Located in East Hanover, NJ Novartis Pharmaceuticals Corporation – an affiliate of Novartis – is reimagining medicine to improve and extend people's lives. As a leading global medicines company, we use innovative science and digital technologies to create transformative treatments in areas of great medical need. In our quest to find new medicines, we consistently rank among the world's top companies investing in research and development. Novartis employs nearly 16,000 people in the United States. For more information, please visit <https://www.novartis.us>.

Novartis and Novartis US is on Twitter. Sign up to follow @Novartis at <https://twitter.com/novartisnews> and @NovartisUS at <https://twitter.com/NovartisUS>.

For Novartis multimedia content, please visit <https://www.novartis.com/news/media-library>.

For questions about the site or required registration, please contact media.relations@novartis.com.

#

Novartis Media Relations

E-mail: media.relations@novartis.com

Julie Masow

Jamie Bennett

Head, US External Engagement Director, US External Engagement

+1 862 579 8456

+1 862 217 3976

julie.masow@novartis.com

jamie.bennett@novartis.com

North America

Sloan Simpson +1 862 778 5052

References

1. ENTRESTO [prescribing information]. East Hanover, NJ: Novartis Pharmaceuticals Corp; February 2021.
2. Yancy C, Jessup M, Bozkurt B. 2013 ACCF/AHA guideline for the management of heart failure: a report of the American College of Cardiology Foundation/American Heart Association task force on practice guidelines. *Circulation*. 2013;128:e240–e327. <https://doi.org/10.1161/CIR.0b013e31829e8776>
3. Fonarow G, Stough W, Abraham W, et al. Characteristics, treatments, and outcomes of patients with preserved systolic function hospitalized for heart failure: a report from the OPTIMIZE-HF registry. *J Am Coll Cardiol*. 2007;50:768-777. doi:10.1016/j.jacc.2007.04.064
4. Virani S, Alonso A, Benjamin E, et al. Heart disease and stroke statistics-2020 update: a report from the American Heart Association. *Circulation*. 2020;141:e139-e596. doi:10.1161/CIR.0000000000000757
5. Solomon S, Rizkala A, Gong J, et al. Angiotensin receptor neprilysin inhibition in heart failure with preserved ejection fraction: rationale and design of the PARAGON-HF trial. *JACC Heart Fail*. 2017;5(7):471-482. doi:10.1016/j.jchf.2017.04.013
6. Solomon S, McMurray J, Anand I, et al. Angiotensin-neprilysin in heart failure with preserved ejection fraction. *N Engl J Med*. 2019;381:1609-1620. doi:10.1056/NEJMoa1908655
7. Gheorghiade M, De Luca L, Fonarow G, et al. Pathophysiologic targets in the early phase of acute heart failure syndromes. *Am J Cardiol*. 2005;96(6A):11G-17G.
8. Dharmarajan K, Hsieh A, Lin Z, et al. Diagnoses and timing of 30-day readmissions after hospitalization for heart failure, acute myocardial infarction, or pneumonia. *JAMA*. 2013;309(4):355-363. doi:10.1001/jama.2012.216476
9. Bueno H, Ross J, Wang Y, et al. Trends in length of stay and short-term outcomes among Medicare patients hospitalized for heart failure: 1993-2008. *JAMA*. 2010;303(21):2141-2147. doi:10.1001/jama.2010.748
10. Langenickel T, Dole W. Angiotensin receptor-neprilysin inhibition with LCZ696: a novel approach for the treatment of heart failure. *Drug Discov Today*. 2012;9(4):e131-e139. doi:10.1016/j.ddstr.2013.11.002

Disclaimer

This press release contains forward-looking statements within the meaning of the United States Private Securities Litigation Reform Act of 1995. Forward-looking statements can generally be identified by words such as "potential," "can," "will," "may," "could," "would," "commitment," "to reduce," "increasing," "goal," "to generate," or similar terms, or by express or implied discussions regarding potential marketing approvals, new indications or labeling for Entresto, or regarding potential future revenues from Entresto. You should not place undue reliance on these statements. Such forward-looking statements are based on our current beliefs and expectations regarding future events, and are subject to significant known and unknown risks and uncertainties. Should one or more of these risks or uncertainties materialize, or should underlying assumptions prove incorrect, actual results may vary materially from those set forth in the forward-looking statements. There can be no guarantee that Entresto will be submitted or approved for sale or for any additional indications or labeling in any market, or at any particular time. Nor can there be any guarantee that Entresto will be commercially successful in the future. In particular, our expectations regarding Entresto could be affected by, among other things, the uncertainties inherent in research and development, including clinical trial results and additional analysis of existing clinical data; regulatory actions or delays or government regulation generally;

global trends toward health care cost containment, including government, payor and general public pricing and reimbursement pressures and requirements for increased pricing transparency; our ability to obtain or maintain proprietary intellectual property protection; the particular prescribing preferences of physicians and patients; general political, economic and business conditions, including the effects of and efforts to mitigate pandemic diseases such as COVID-19; safety, quality, data integrity or manufacturing issues; potential or actual data security and data privacy breaches, or disruptions of our information technology systems, and other risks and factors referred to in Novartis AG's current Form 20-F on file with the US Securities and Exchange Commission. Novartis is providing the information in this press release as of this date and does not undertake any obligation to update any forward-looking statements contained in this press release as a result of new information, future events or otherwise.

Source URL: <https://www.novartis.com/us-en/news/media-releases/novartis-entresto-granted-expanded-indication-chronic-heart-failure-fda>

List of links present in page

- <https://www.novartis.com/us-en/us-en/news/media-releases/novartis-entresto-granted-expanded-indication-chronic-heart-failure-fda>
- <https://www.multivu.com/players/English/8848351-novartis-entresto-fda-approval/>
- https://www.novartis.com/us-en/us-en/sites/novartis_us/files/2022-03/entresto_0.pdf
- <http://www.fda.gov/medwatch>
- <http://www.entresto.com/>
- <https://www.novartis.com/us-en/us-en/home>
- <https://twitter.com/novartisnews>
- <https://twitter.com/NovartisUS>
- <https://www.novartis.com/news/media-library>
- <mailto:media.relations@novartis.com>
- <mailto:julie.masow@novartis.com>
- <mailto:jamie.bennett@novartis.com>
- <mailto:investor.relations@novartis.com>
- <https://doi.org/10.1161/CIR.0b013e31829e8776>